

## XANTARES BIOCIENTÍFICOS

Desde o grupo de traballo de formación e divulgación de **BioReDes** queremos propoñer un **espazo regular de interacción** para dar visibilidade a **temas científicos** de especial interese polo seu impacto social, económico e ambiental, así como polo seu grao de innovación. Para iso, celebraremos o primeiro venres de cada mes (o segundo venres se coincide con pontes ou festivos) un seminario científico que bautizamos como **Xantares BioCientíficos**. Consistirán en **charlas curtas** (30 minutos aprox.), impartidas por **investigadores invitados ou da propia agrupación**, presentadas nun **ton divulgativo** (búscase que sexan accesibles para investigadores non especialistas no tema) e seguidas dun **debate aberto tanto científico como de impacto social**. Pensamos acompañalo dun xantar lixeiro ao finalizar porque pretendemos que esta actividade se converta nun espazo informal para coñecernos mellor e para favorecer o **intercambio de ideas** e promover **futuras colaboracións**. Ademais, estes encontros poden ser un excelente complemento para a **formación dos investigadores máis novos** (acreditarase a asistencia), polo que vos pedimos que animedes a participar aos doutorandos dos vosos grupos.

A primeira cita será o **venres 8 de marzo ás 13:00 horas no edificio CACTUS** cunha charla titulada **“Globalización e patoloxías na produción de moluscos: cancro contaxioso e marteiliosis”**, impartida polo Dr. Antonio Villalba, investigador do Centro de Investigacións Mariñas e coordinador da área de Patoloxía. Para que poidades ter información de antemán sobre a importancia do tema achegámosvos un artigo científico e un breve resumo da presentación.

Aproveitamos para comunicarvos que, se tedes suxestións sobre charlas de interese para os Xantares BioCientíficos, podedes remitilas a Iago Rodríguez, técnico de I+D da Agrupación ([iago.palmeiro@usc.es](mailto:iago.palmeiro@usc.es))

Esperámoste! Ven a tomar un petisco con BioReDes!

### Grupo de traballo de formación e divulgación

Dra. Belén Díaz (AGRONOMÍA)

Dr. Gonzalo Fernández (INVESAGA)

Dra. Azucena Mora (LREC)

Dr. Juan Ortiz (CIGEO)

Dra. Maribel Quiroga (GAPAVET, coordinadora do grupo)

Dr. Paulino Martínez, (ACUIGEN, director de BioReDes)

## Breve resumo da presentación

### “Globalización e patoloxías na produción de moluscos: Cancro contaxioso e marteiliosis”

Antonio Villalba, David Iglesias, Asunción Cao, María Jesús Carballal

*Centro de Investigacións Mariñas, Consellería do Mar, Xunta de Galicia*

A produción de moluscos, derivada de acuicultura e pesca, contribúe de forma crecente á alimentación humana. Como noutros casos de produción animal, algunhas enfermidades ameazan de forma dramática as explotacións marisqueiras. En Galicia a produción marisqueira ten especial relevancia e, por iso, algunhas enfermidades de moluscos supoñen unha **convulsión no ámbito socioeconómico**; o caso máis recente é a marteiliosis do berberecho.

A investigación desenvolvida ao redor dun tipo de cancro de moluscos que lembra á leucemia conduciu a un achado sorprendente que **convulsionou o mundo científico**: o contaxio directo do cancro entre individuos dunha mesma especie e mesmo de especies distintas mediante o tránsito de células cancerosas liberadas á auga circundante por individuos enfermos e que se introducen en individuos sans, proliferando nestes. A evidencia de que a transmisión de cancro no medio acuático non é excepcional está a deparar novas perspectivas sobre o cancro.

# Widespread transmission of independent cancer lineages within multiple bivalve species

Michael J. Metzger<sup>1,2</sup>, Antonio Villalba<sup>3,4</sup>, María J. Carballal<sup>3</sup>, David Iglesias<sup>3</sup>, James Sherry<sup>5</sup>, Carol Reinisch<sup>5</sup>, Annette F. Muttray<sup>6,7</sup>, Susan A. Baldwin<sup>6</sup> & Stephen P. Goff<sup>1,2,8</sup>

**Most cancers arise from oncogenic changes in the genomes of somatic cells, and while the cells may migrate by metastasis, they remain within that single individual. Natural transmission of cancer cells from one individual to another has been observed in two distinct cases in mammals (Tasmanian devils<sup>1</sup> and dogs<sup>2,3</sup>), but these are generally considered to be rare exceptions in nature. The discovery of transmissible cancer in soft-shell clams (*Mya arenaria*)<sup>4</sup> suggested that this phenomenon might be more widespread. Here we analyse disseminated neoplasia in mussels (*Mytilus trossulus*), cockles (*Cerastoderma edule*), and golden carpet shell clams (*Polititapes aureus*) and find that neoplasias in all three species are attributable to independent transmissible cancer lineages. In mussels and cockles, the cancer lineages are derived from their respective host species; however, unexpectedly, cancer cells in *P. aureus* are all derived from *Venerupis corrugata*, a different species living in the same geographical area. No cases of disseminated neoplasia have thus far been found in *V. corrugata* from the same region. These findings show that transmission of cancer cells in the marine environment is common in multiple species, that it has originated many times, and that while most transmissible cancers are found spreading within the species of origin, cross-species transmission of cancer cells can occur.**

Disseminated neoplasia, or haemic neoplasia, a leukaemia-like disease, occurs with high prevalence in multiple bivalve species<sup>5,6</sup>. Here we investigate the possibility that cancers in three species could be attributed to transmissible cancer cells, and whether these cancers are restricted to the species of origin or can undergo cross-species transmission.

Mussels (*M. trossulus*; Fig. 1a) are subject to disseminated neoplasia in the Pacific Northwest Coast<sup>7,8</sup>, and evidence of common polymorphisms in neoplasias suggested that these might represent a transmissible cancer<sup>9</sup>. Twenty-eight mussels (*M. trossulus*) collected from West Vancouver were screened for neoplasia by drawing haemolymph and analysing haemocytes for the rounded, non-adherent morphology of neoplastic cells. Two were identified with high levels of neoplastic cells. We sequenced part of the mitochondrial cytochrome *c* oxidase I (*mtCOI*) gene in host tissue and neoplastic haemocytes from the two diseased animals and four normal animals to test whether the neoplastic cells were derived from the host individuals or exhibited a distinct genotype, the hallmark of transmissible neoplasia. While solid tissue and haemocyte genotypes within each normal animal were always identical, solid tissue and neoplastic haemocyte genotypes of the diseased animals were discordant (Fig. 1b and Extended Data Fig. 1). Moreover, the same single nucleotide polymorphisms (SNPs) were found in neoplastic cells of the two diseased individuals, indicating that the cancer cells were not of host origin and suggesting that they arose from a single clonal origin. *EF1 $\alpha$*  gene sequences also revealed that the genotypes of the host cells and neoplastic haemocytes were discordant, and that the

genotypes of the neoplastic cells of the two different animals were again identical to each other (Fig. 1c).


To determine whether this transmissible cell line was widespread in the *M. trossulus* population, 250 were collected from Vancouver Island, and seven potentially diseased individuals were analysed. In one neoplastic sample (MW81), the haemocyte and tissue genotypes did not match, and the haemocytes contained the same *mtCOI* allele and the same *EF1 $\alpha$*  major and minor alleles found in the other samples (Fig. 1b, c and Extended Data Fig. 1). These genotypes strongly indicate the existence of a *M. trossulus*-derived transmissible cancer lineage circulating in the wild population.

High prevalence of disseminated neoplasia has been observed in two species of bivalves on the Galician Coast: cockles (*C. edule*)<sup>10,11</sup> and golden carpet shell clams (*P. aureus*, previously named *Venerupis aurea*)<sup>12</sup>. The disease in cockles (*C. edule*) exhibits one of two distinct morphologies, termed types A and B<sup>13,14</sup>. We collected about 150 cockles (*C. edule*; Fig. 2a), and examined the genotypes of solid tissue and haemocytes of six normal individuals and six with high (>75%) or moderate (15–75%) amounts of neoplastic cells in the haemolymph. Nine polymorphic microsatellite loci<sup>15</sup> were amplified from normal animals, and allele sizes from tissue and haemocytes of each normal animal all matched, but allele sizes in haemocytes and tissue of diseased animals were discordant (Fig. 2b, c). In a phylogenetic tree based on microsatellite alleles<sup>16,17</sup> the neoplastic haemocyte genotypes did not group with the host tissue genotypes, consistent with transmissible cancer, and instead clustered into two distinct branches, suggesting two independent cancer lineages (Fig. 2d). We sequenced the *mtCOI* gene and identified several SNPs that were present only in lineage 2, and not in any of the normal animals. No unique SNPs were identified in the *mtCOI* region sequenced in lineage 1.

The microsatellite alleles and *mtCOI* SNPs suggest two independent cancer lineages, but these data are also consistent with two subgroups that have diverged from a single transmissible cancer lineage. To investigate this question, we sequenced an approximately 3-kilobase intron-spanning region in *EF1 $\alpha$*  from six normal individuals and two diseased individuals from each lineage (Fig. 2e). Both neoplastic haemocyte alleles were different from the tissue alleles in all diseased individuals, and the two alleles of the neoplasm in each diseased individual were nearly identical to the two alleles of the other individual in the lineage. However, both alleles in lineage 1 cells were different from those in lineage 2. Moreover, the neoplastic alleles were more closely related to some normal alleles than they were to the alleles of the alternative lineage. These data strongly suggest an independent origin of these two lineages.

Histological and morphometric examination showed that all three samples in lineage 1 were type A (characterized by a looser arrangement of neoplastic cells in the connective tissue, with pleomorphic nuclei) and all three samples in lineage 2 were type B (tighter arrangement,

<sup>1</sup>Department of Biochemistry and Molecular Biophysics, Columbia University, New York 10032, USA. <sup>2</sup>Howard Hughes Medical Institute, New York, New York 10032, USA. <sup>3</sup>Centro de Investigacións Mariñas, Consellería do Mar, Xunta de Galicia, Vilanova de Arousa 36620, Spain. <sup>4</sup>Department of Life Sciences, University of Alcalá, Alcalá de Henares 28871, Spain. <sup>5</sup>Environment Canada, Water Science & Technology Directorate, Burlington, Ontario L7R 4A6, Canada. <sup>6</sup>Chemical and Biological Engineering, University of British Columbia, Vancouver V6T 1Z3, Canada. <sup>7</sup>SLR Consulting Canada Ltd., Vancouver V6J 1V4, Canada. <sup>8</sup>Department of Microbiology and Immunology, Columbia University, New York, New York 10032, USA.



**Figure 1 | Analysis of tissue and haemocyte genotypes of normal mussels (*M. trossulus*) and mussels with disseminated neoplasia using mitochondrial and nuclear DNA markers. a, Representative *M. trossulus*, with ruler for scale. b, Sequencing of *mtCOI* (bases 99–759) was conducted for both host solid tissue (T) and haemocytes (H) of normal ( $n = 4$ ) and neoplastic mussels ( $n = 3$ ). Open boxes mark SNPs that differ from allele A. Filled boxes mark discordance between neoplastic haemocyte and tissue genotypes. Sequences are numbered using AY823625.1. c, Maximum likelihood phylogenetic tree based on sequences of an intron-containing**

region of *EF1α*, with bootstrap values over 50 shown and a scale bar showing genetic distance. *M. edulis* (614 base pairs (bp), EU684203.1) was used as an outgroup. In normal animals, the sequences of tissue and haemocyte DNA were identical and are presented together (T&H; black circles). For neoplastic animals, the tissue (T; open red circles) and haemocyte (H; filled red circles) alleles differed. Letters 'a' and 'b' denote multiple alleles in heterozygous individuals, and major and minor alleles from neoplastic alleles are marked. See Extended Data Fig. 1 for further details.

with smaller cells than type A and rounded, smaller nuclei; Fig. 2f–i and Extended Data Table 1). Altogether, these results argue that two distinct lineages of transmissible cancer, with distinct morphologies and genotypes, arose independently in cockles and are circulating in this species.

We also examined golden carpet shell clams (*P. aureus*; Fig. 3a), which are present in the same habitat as several other bivalves, including the closely related pullet shell clam (*V. corrugata*; Fig. 3b). Of 74 *P. aureus* individuals tested, 9 had high levels of disease and 22 had low-to-medium disease. We sequenced regions of *mtCOI*, the ribosomal DNA internal transcribed spacer (rDNA *ITS*), and *EF1α* from tissue and haemocyte DNA of six highly diseased and six normal *P. aureus* individuals, and again found that the genotypes of neoplastic cells were nearly identical and did not match those of their hosts (Fig. 3c–e). In contrast to the transmissible cancers in other species, however, the sequences of the neoplastic cells were highly dissimilar from normal host sequence (only 78.4–78.5% identical in the *mtCOI* locus and 89.3–93.2% identical at nuclear loci, ignoring insertions and deletions). The neoplastic genotypes were instead near-perfect matches to the sequences of *V. corrugata* (98.6–7% and 99.3–100% identical, respectively).

To confirm that detection of the *V. corrugata* sequence reflected the neoplastic cells observed morphologically, we used species-specific *EF1α* quantitative PCR (qPCR) to quantify the fraction of cells that were derived from each species in the tissue and haemocyte samples (Fig. 3f). As expected, no *V. corrugata* sequences were detected in normal *P. aureus* animals, but high frequencies of *V. corrugata* DNA were detected in all six individuals diagnosed with high levels of neoplasia and lower amounts in most individuals diagnosed with medium and low levels. The small number of individuals diagnosed with low and medium disease in which *V. corrugata* DNA was not detected could have primary host-derived neoplasia, a different transmissible cancer lineage, too low an abundance of tumour DNA to detect, or even a small amount of normal cells with an unusually rounded morphology. In diseased animals, the highest levels of *V. corrugata* sequence were detected in haemocytes, and lower levels were present in tissues, probably because of infiltration of cancer cells via the circulation. We conclude that the cancer spreading in the *P. aureus* population resulted


from cross-species transmission of cancer cells of *V. corrugata* origin. It is noteworthy that no cases of disseminated neoplasia have thus far been found in *V. corrugata* from the same region, despite analysis of hundreds of clams, suggesting that the *V. corrugata* cancer cells do not initiate disease in the species of origin and were only observed to colonize *P. aureus* animals.

We previously identified an LTR-retrotransposon, *Steamer*, which was amplified in the transmissible neoplasia lineage in the soft-shell clam (*M. arenaria*)<sup>18</sup>. Using degenerate primers, we identified at least one *Steamer*-like element (*SLE*) in each of the three species studied here. In each case, copy numbers were variable among individuals, but no massive amplification of these particular retrotransposons was observed in any of the transmissible cancers assayed (Extended Data Fig. 2). This suggests that *Steamer*-like retrotransposon amplification is not essential in development of transmissible clones.

Our results indicate that transmission of contagious cancer cells is a widespread phenomenon in the marine environment, with multiple independent lineages developing in multiple species in four bivalve families. Along with the recent identification of a second independent lineage of transmissible cancer in Tasmanian devils<sup>19</sup>, these findings confirm that, under suitable conditions, the development of transmissible cancer can occur multiple times, and suggests that some species may be more susceptible to development of transmissible neoplasia than others.

Spontaneous haemic neoplasia must occur at some frequency for transmissible lineages to arise, but cases of transmissible cancer appear to outnumber spontaneous disease, at least in the species investigated so far (that is, all neoplastic cases tested in cockles, golden carpet shell clams, and three of nine samples in mussels, in addition to all samples in the previous report of soft-shell clams<sup>4</sup>).

Transmissible cancers appear largely restricted to the species of origin in nature. While canine transmissible venereal tumour has been experimentally transplanted to coyotes<sup>20</sup>, jackals<sup>21</sup>, and foxes<sup>22–24</sup>, no examples of natural transmission beyond dogs have been reported<sup>25</sup>. Disseminated neoplasias of molluscs have only been transplanted to members of the same species (soft-shell clams, mussels, and others<sup>5,6</sup>). Attempts to transfer *M. arenaria* neoplasia through water exposure to both *M. arenaria* and *M. trossulus* only resulted in transfer




**Figure 2 | Analysis of lineages of transmissible neoplasia in cockles (*C. edule*).** **a**, Representative *C. edule* (scale bar, 20 mm). **b**, **c**, Microsatellite loci were amplified from solid tissue (T) and haemocyte (H) DNA from normal (samples N1–N6) and diseased (M1–M3 and H1–H3) cockles. Products of a representative locus (CeATC1–5) are shown. Triangles mark samples corresponding to lineages 1 (red) and 2 (purple). **d**, Neighbour-joining phylogenetic tree based on nine microsatellite loci (see Source data). Bootstrap values over 50 are shown and scale bar is

based on comparison across all 104 observed alleles. *mtCOI* SNPs unique to lineage 2 are marked. **e**, Maximum likelihood tree of *EF1α* (2725–4249 bp, spanning four introns), rooted at the midpoint, with bootstrap values above 50 and scale bar showing genetic distance. Letters 'a' and 'b' denote multiple alleles from heterozygous samples, and alleles from neoplastic lineages 1 and 2 are marked. **f**, **g**, Histology of cockles H3 and H2, representative of type A neoplastic cells. **h**, **i**, Histology of cockle H1, representative of type B neoplastic cells.

of disease to *M. arenaria*<sup>26</sup>, and injection of *M. trossulus* cancer cells into multiple bivalve species only resulted in engraftment in *M. trossulus*<sup>27</sup>. Our finding that multiple cancer lineages are most often found to spread within the original host species is consistent with these previous experiments, and suggests that there may be species-specific restriction factors that prevent engraftment into divergent hosts.

In this context, our observation of cross-species transmission, a cancer from one species spreading through another, is particularly striking. They both belong to the Veneridae family (some studies suggest that *P. aureus* should belong to the *Venerupis* genus<sup>28</sup>), and coexist in the same beds. This close relationship may have aided the transmission.

It is notable that despite ongoing surveillance of the bivalves of the Galician coast since the 1990s, only one *V. corrugata* has been identified as harbouring disseminated neoplasia (S. Darriba, personal communication), while the prevalence of disease in *P. aureus* is quite high<sup>12</sup> (12% of the individuals collected for the current study had high levels of neoplasia and 42% had some detectable level of disease). This would be explained if the cancer originated in *V. corrugata*, led to selective loss of susceptible animals, and left the current population of *V. corrugata* resistant to engraftment and disease. The spread of transmissible cancers and the evolutionary pressure to defend against them may be a strong and underappreciated selective force in the evolution of multicellular organisms.



**Figure 3 | Phylogenetic analysis of neoplastic cells in *P. aureus* and quantification of *V. corrugata* cell engraftment in normal and diseased animals. a, Representative *P. aureus* clams (scale bar, 20 mm). b, Representative *V. corrugata* clams (scale bar, 20 mm). c–e, Maximum likelihood trees of sequences from solid tissue (T) and haemocyte (H) DNA from normal (N1–N6) and highly diseased (H1–H6) *P. aureus* and normal *V. corrugata*, based on (c) *mtCOI* (658 bp), (d) *rDNA ITS* (454–373 bp), and (e) *EF1α* (148 bp, letters ‘a’ and ‘b’ denote multiple**

alleles). Open black circles, tissue and haemocytes of normal *P. aureus*; open red circles, tissue of diseased *P. aureus*; filled red circles, haemocytes of diseased *P. aureus*; scale bars show genetic distance. f, Quantification of cancer cell engraftment using species-specific amplification of *EF1α* from haemocyte and tissue DNA of normal *V. corrugata*, normal *P. aureus*, and *P. aureus* diagnosed as having low, moderate, or high levels of disseminated neoplasia.

In summary, our findings show that transmission of cancers between individuals within a species may be more common than previously assumed, and that transmission can even occur between species. We now know of eight transmissible cancers in nature: one lineage in dogs, two lineages in Tasmanian devils, and five lineages circulating in four species of molluscs, including one example of cross-species transmission. Further studies may well reveal additional examples. The recent report of a cancer composed of transformed tapeworm cells in an immunocompromised AIDS patient<sup>29</sup> highlights the possibility that transmissible tumours could arise in humans. These transmissible cancers constitute a distinct class of infectious agent and show the remarkable ability of tumours to acquire new phenotypes that promote their own survival and propagation.

**Online Content** Methods, along with any additional Extended Data display items and Source Data, are available in the online version of the paper; references unique to these sections appear only in the online paper.

Received 29 January; accepted 24 May 2016.

Published online 22 June 2016.

1. Pearse, A. M. & Swift, K. Allograft theory: transmission of devil facial-tumour disease. *Nature* **439**, 549 (2006).
2. Murgia, C., Pritchard, J. K., Kim, S. Y., Fassati, A. & Weiss, R. A. Clonal origin and evolution of a transmissible cancer. *Cell* **126**, 477–487 (2006).

3. Rebbeck, C. A., Thomas, R., Breen, M., Leroi, A. M. & Burt, A. Origins and evolution of a transmissible cancer. *Evolution* **63**, 2340–2349 (2009).
4. Metzger, M. J., Reinisch, C., Sherry, J. & Goff, S. P. Horizontal transmission of clonal cancer cells causes leukemia in soft-shell clams. *Cell* **161**, 255–263 (2015).
5. Barber, B. J. Neoplastic diseases of commercially important marine bivalves. *Aquat. Living Resour.* **17**, 449–466 (2004).
6. Carballal, M. J., Barber, B. J., Iglesias, D. & Villalba, A. Neoplastic diseases of marine bivalves. *J. Invertebr. Pathol.* **131**, 83–106 (2015).
7. Moore, J. D., Elston, R. A., Drum, A. S. & Wilkinson, M. T. Alternate pathogenesis of systemic neoplasia in the bivalve mollusc *Mytilus*. *J. Invertebr. Pathol.* **58**, 231–243 (1991).
8. Vassilenko, E. & Baldwin, S. A. Using flow cytometry to detect haemic neoplasia in mussels (*Mytilus trossulus*) from the Pacific Coast of Southern British Columbia, Canada. *J. Invertebr. Pathol.* **117**, 68–72 (2014).
9. Vassilenko, E. I., Muttray, A. F., Schulte, P. M. & Baldwin, S. A. Variations in p53-like cDNA sequence are correlated with mussel haemic neoplasia: a potential molecular-level tool for biomonitoring. *Mutat. Res.* **701**, 145–152 (2010).
10. Villalba, A., Carballal, M. J. & López, C. Disseminated neoplasia and large foci indicating heavy haemocytic infiltration in cockles *Cerastoderma edule* from Galicia (NW Spain). *Dis. Aquat. Organ.* **46**, 213–216 (2001).
11. Díaz, S., Iglesias, D., Villalba, A. & Carballal, M. J. Long-term epidemiological study of disseminated neoplasia of cockles in Galicia (NW Spain): temporal patterns at individual and population levels, influence of environmental and cockle-based factors and lethality. *J. Fish Dis.* <http://dx.doi.org/10.1111/jfd.12436> (2016).
12. Carballal, M. J., Iglesias, D., Díaz, S. & Villalba, A. Disseminated neoplasia in clams *Venerupis aurea* from Galicia (NW Spain): histopathology, ultrastructure and ploidy of the neoplastic cells, and comparison of diagnostic procedures. *J. Invertebr. Pathol.* **112**, 16–19 (2013).

13. Carballal, M. J., Iglesias, D., Santamarina, J., Ferro-Soto, B. & Villalba, A. Parasites and pathologic conditions of the cockle *Cerastoderma edule* populations of the coast of Galicia (NW Spain). *J. Invertebr. Pathol.* **78**, 87–97 (2001).
14. Iglesias, D. *Estudio Patológico de las Poblaciones de Berberecho Cerastoderma edule (L.) de Galicia*. Ph.D. thesis, Univ. Santiago do Compostela (2006).
15. Martínez, L., Arias, A., Mendez, J., Insua, A. & Freire, R. Development of twelve polymorphic microsatellite markers in the edible cockle *Cerastoderma edule* (Bivalvia: Cardiidae). *Conserv. Genet. Resour.* **1**, 107–109 (2009).
16. Kamvar, Z. N., Tabima, J. F. & Grünwald, N. J. Poppr: an R package for genetic analysis of populations with clonal, partially clonal, and/or sexual reproduction. *PeerJ* **2**, e281 (2014).
17. Rodzen, J. A. & May, B. Inheritance of microsatellite loci in the white sturgeon (*Acipenser transmontanus*). *Genome* **45**, 1064–1076 (2002).
18. Arriagada, G. *et al.* Activation of transcription and retrotransposition of a novel retroelement, Steamer, in neoplastic hemocytes of the mollusk *Mya arenaria*. *Proc. Natl Acad. Sci. USA* **111**, 14175–14180 (2014).
19. Pye, R. J. *et al.* A second transmissible cancer in Tasmanian devils. *Proc. Natl Acad. Sci. USA* **113**, 374–379 (2016).
20. Cockrill, J. M. & Beasley, J. N. Transmission of transmissible venereal tumor of the dog to the coyote. *Am. J. Vet. Res.* **40**, 409–410 (1979).
21. Samsó, A. *Recherches Experimentales sur le Sarcome de Sticker*. Docteur ès Sciences Naturales thesis, Univ. Paris (1965).
22. Sticker, A. Transplantables Rundzellensarkom des Hundes. *Z. Krebsforsch.* **4**, 227–314 (1906).
23. Wade, H. An experimental investigation of infective sarcoma of the dog, with a consideration of its relationship to cancer. *J. Pathol. Bacteriol.* **12**, 384–425 (1908).
24. Dungen, V. Zur Biologie des Rundzellensarkoms des Hundes. *Munch. med. Wochenschr.* No. 5, 238–239 (1912).
25. Strakova, A. & Murchison, E. P. The changing global distribution and prevalence of canine transmissible venereal tumour. *BMC Vet. Res.* **10**, 168 (2014).
26. Mateo, D. R., MacCallum, G. S. & Davidson, J. Field and laboratory transmission studies of haemic neoplasia in the soft-shell clam, *Mya arenaria*, from Atlantic Canada. *J. Fish Dis.* <http://dx.doi.org/10.1111/jfd.12426> (2015).
27. Kent, M. L., Wilkinson, M. T., Drum, A. S. & Elston, R. A. Failure of transmission of hemic neoplasia of bay mussels, *Mytilus trossulus*, to other bivalve species. *J. Invertebr. Pathol.* **57**, 435–436 (1991).
28. Canapa, A., Schiaparelli, S., Marota, I. & Barucca, M. Molecular data from the 16S rRNA gene for the phylogeny of Veneridae (Mollusca: Bivalvia). *Mar. Biol.* **142**, 1125–1130 (2003).
29. Muehlenbachs, A. *et al.* Malignant transformation of *Hymenolepis nana* in a human host. *N. Engl. J. Med.* **373**, 1845–1852 (2015).

**Acknowledgements** M.J.M. and S.P.G. were supported by the Howard Hughes Medical Institute and Training Grant T32 CA009503. D.I., M.J.C., and A.V. were supported by the Consellería do Mar da Xunta de Galicia, through the project PGIDIT-CIMA 13/03. We thank J. Ausió for help in collection of *M. trossulus* from Vancouver Island.

**Author Contributions** M.J.M. and S.P.G. wrote the manuscript. M.J.M. conducted molecular analyses. A.F.M. and S.A.B. collected and diagnosed *M. trossulus* from West Vancouver. J.S. and C.R. collected and diagnosed *M. trossulus* from Vancouver Island. D.I., M.J.C., and A.V. collected and diagnosed *C. edule* and *P. aureus*. M.J.C. produced micrographs of *C. edule* neoplastic haemocytes, and D.I. conducted morphometric analysis.

**Author Information** Sequences generated in this work have been deposited in GenBank under accession numbers KX018521–KX018605. Reprints and permissions information is available at [www.nature.com/reprints](http://www.nature.com/reprints). The authors declare no competing financial interests. Readers are welcome to comment on the online version of the paper. Correspondence and requests for materials should be addressed to S.P.G. (spg1@cumc.columbia.edu).

**Reviewer Information** *Nature* thanks E. Murchison, S. O'Brien, G. De Vico, R. A. Weiss and the other anonymous reviewer(s) for their contribution to the peer review of this work.

## METHODS

No statistical methods were used to predetermine sample size. The experiments were not randomized. The investigators were not blinded to diagnosis during sequencing and morphology assessment.

**Collection and diagnosis of *M. trossulus*.** Mussel (*M. trossulus*) specimens 5–6 cm long were collected from the intertidal zone at low tide (noon) on 18 April 2015 at Copper Beach (49° 22' 41" N, 123° 16' 44" W, West Vancouver, British Columbia, Canada). They were transported to the laboratory in aerated seawater from Copper Beach. In the laboratory, 0.5–1.0 ml haemolymph was removed from the posterior adductor muscle. For each individual, one drop of haemolymph was placed on a poly-L-lysine-coated slide and let sit for 10–15 min to allow the cells to spread and attach to the slide. Thereafter, the slide was viewed under a Zeiss Axiostar light microscope at  $\times 40$  magnification. Normal (non-neoplastic) specimens were those with greater than 90% cells with normal appearance: that is, agranular or granular haemocytes with spread pseudopodia<sup>30</sup>. Fully leukaemic (diseased) specimens were those with prolific amounts (>90%) of round, non-adherent cells. Haemolymph was added to 1.5 ml Eppendorf tubes and spun at 900g for 3 min. After the supernatant was withdrawn, the cells were re-suspended in absolute ethanol before shipping. Excised tissues (mantle, foot, and gills) were preserved in absolute ethanol before shipping. Of 28 individuals collected from West Vancouver, two had high levels of neoplastic disease.

A second set of *M. trossulus* samples were collected from Esquimalt, Vancouver Island, British Columbia, Canada. Haemolymph was extracted and cell morphology was used for diagnosis as above. Haemocyte and solid tissue samples were fixed in ethanol before DNA extraction. Of 250 individuals collected from Esquimalt, 9 were scored as potentially moderately or highly diseased, with 7 samples available for analysis.

**Collection and diagnosis of *C. edule* and *P. aureus*.** Cockles (*C. edule*) were collected from an intertidal bed named O Sarrido (42° 30' N, 8° 49' W) and golden carpet shell clams (*P. aureus*) and pullet shell clams (*V. corrugata*) were collected from a subtidal bed named O Bohido (42° 32' N, 8° 51' W); both shellfish beds are located in the ria of Arousa (Galicia, northwest Spain). Once in the laboratory of Centro de Investigaciones Mariñas, each cockle and clam was notched through the shell margin close to the posterior adductor muscle and haemolymph (as much as possible) was collected from the posterior adductor muscle using a 2-ml syringe with a 21-gauge needle. A small quantity (about 100  $\mu$ l) of haemolymph was used to produce a cell monolayer onto a slide by cytocentrifugation (92g, 5 min, 4°C), which was fixed, stained with a Hemacolor (Merck) kit, and examined with light microscopy for diagnosis of disseminated neoplasia; the remaining haemolymph was preserved in absolute ethanol for molecular analysis. After haemolymph collection, molluscs were shucked and a small piece of mantle (about 5 mm  $\times$  5 mm) was removed and preserved in absolute ethanol for molecular analysis; additionally, an approximately 5-mm-thick section of meat, containing gills, visceral mass, foot, and mantle lobes, was fixed in Davidson's solution and embedded in paraffin. Sections of 5  $\mu$ m thickness were stained with Harris's haematoxylin and eosin<sup>31</sup>. Histological sections were examined under light microscopy for histopathological analysis.

Morphometric analysis was conducted on histological sections of cockle (*C. edule*) samples. Neoplastic cells (both A and B types) had a unique morphology, and were clearly distinguished from normal cells, with much larger overall size, and much larger nucleus. The longest diameters of the cell and the nucleus of at least ten neoplastic cells for each individual were measured by direct examination of histological sections with light microscopy using a reticle. Two-tailed *t*-tests were used for comparisons of morphometric data from different individuals and types.

Cockles (*C. edule*) and golden carpet shell clams (*P. aureus*) were ranked according to a scale of disease severity based on haemolymph diagnosis: non-affected (N0, or N); low severity (N1, or L), when individuals showed proportions of neoplastic cells lower than 15% in the haemolymph cell monolayers; moderate severity (N2, or M), when the proportion ranged from 15% to 75%; and high severity (N3, or H), when the proportion was higher than 75% (ref. 32). Seventy-four golden carpet shell clams (*P. aureus*) were collected with 12 diagnosed with light, 10 with moderate, and 9 with heavy neoplasia. About 150 cockles (*C. edule*) were collected and a subset was analysed both for disease and for morphological type of neoplastic cells. Of the 30 in this subset, two had type A neoplastic cells (one light and the other moderate) and one had low levels of type B neoplastic cells. We have also collected and analysed hundreds of pullet clams (*V. corrugata*) from multiple beds throughout Galicia from 1988 to the present, including about 100 from the same bed in which the samples of *P. aureus* were collected for this study (O Bohido). So far, we have not observed any *V. corrugata* samples with neoplastic disease. One *V. corrugata* individual from a different location has been identified by a different researcher as harbouring disseminated neoplasia, with a different morphology than that observed in *P. aureus* neoplasia (S. Darriba, personal communication).

**DNA extraction.** DNA was extracted from ethanol-fixed haemocytes using DNeasy Blood and Tissue Kit (Qiagen). DNA extraction of tissues used the same kit, but included an additional step to reduce the amount of PCR-inhibiting polysaccharides. After tissue lysis, 63  $\mu$ l of buffer P3 was added to lysate and allowed to precipitate for 5 min. Lysate was spun 10 min at full speed at 4°C, and the resulting supernatant was mixed with buffer AL for 10 min at 37°C, then mixed with ethanol and added to the column, continuing with the standard protocol.

***mtCOI*, *EF1 $\alpha$* , *ITS*, Steamer-like PCR.** Primers and annealing temperatures used are listed in Extended Data Table 2. PfuUltra II Fusion HS DNA Polymerase (Agilent) was used to amplify 10 ng of genomic DNA for 35 cycles. Extension was at 72°C for 15 s for *mtCOI* for all bivalves<sup>33</sup>. For *P. aureus*, *ITS* was amplified (using primers modified from refs 34, 35) with extension for 30 s. Despite multiple copies of *ITS* in genomic DNA, a single sequence was observed for each normal *P. aureus* and *V. corrugata*, and a single pair of host/neoplasm sequences was observed in each diseased individual. PCR for *EF1 $\alpha$*  followed the same program, with extension for 20 s in mussels (*M. trossulus*) and clams (*P. aureus* and *V. corrugata*) and 1 min 30 s in cockles (*C. edule*). For amplification of Steamer-like elements, a degenerate primer pair was designed to match conserved regions in reverse transcriptase and integrase of the LTR-retrotransposon, Steamer, and 10 ng of DNA was amplified for 35 cycles, annealing at 55°C for 20 s and extending for 1 min at 72°C. PCR products were directly sequenced. When multiple alleles could not be resolved by direct sequencing, PCR products were cloned using the Zero Blunt TOPO Kit (Invitrogen), and at least six colonies were sequenced. Sequences were aligned with Clustal W, with some manual adjustment. Primer-binding regions were excluded from analysis and are not included in the sizes listed as sequenced. Maximum likelihood phylogenetic trees were generated using PhyML 3.0 (ref. 36), using the HKY85 substitution model, with 100 bootstrap replicates, treating gaps in the alignment as missing data. Trees were visualized using FigTree version 1.4, with addition of markers at the branch termini. Each phylogenetic tree based on alignment of sequence at a single locus (Figs 1c, 2e and 3b–d) includes a scale bar that shows genetic distance (0–1) based on the frequency of nucleotide divergence. All sequences are available in GenBank (accession numbers KX018521–KX018605).

**Analysis of microsatellite loci from *C. edule*.** Microsatellites were amplified using primers for 12 loci, reported previously to be polymorphic in cockle (*C. edule*) populations<sup>15</sup>. Of the 12 primer pairs, 9 pairs amplified products in all cockle DNA samples, with 1–2 alleles observed in normal samples and 1–4 neoplastic alleles observed. These were used in all further analyses, with fluorescent modifications on the 5' end of the forward primers as listed (Extended Data Table 2). KOD polymerase (Millipore) was used to prevent ambiguity of untemplated addition of A residues by *Taq* polymerase. Products were run on an agarose gel for visualization. Allele sizes were identified to single-base precision by fragment analysis using fluorescent primers (6-FAM, PET, NED, and VIC) using a 3730xl Genetic Analyzer with the LZ-500 size standard (Applied Biosystems, operated by Genewiz). Peaks were called using Peak Scanner 2.0 (Applied Biosystems), rounding sizes to whole bases, with some manual adjustment to keep alleles of the same size together.

We used the R package *Poppr*<sup>16</sup> to generate distance matrices and phylogenetic trees using Bruvo's method with the infinite alleles model. The value of *c* (repeat size) was set to 0.00001 to calculate distances using the band sharing model, in which alleles are either identical sizes (distance = 0) or non-identical (distance = 1). Each allele was coded as a dominant marker<sup>17</sup>, with a single variable for each allele observed at each position (104 total markers from 9 loci). For each allele (a single size at a single locus), a sample is observed to be present, absent, or the information missing. For example, a sample with sizes 195 and 199 at locus A would be marked as 'present' for marker 'A-195' and 'A-199', but 'absent' for all other sizes at that locus (like 'A-203' and 'A-207'). This analysis method ignores the uncertainty in the copy number of each locus in aneuploid cancer cells and allows for ambiguity at particular alleles. In cases where the haemocyte and tissue genotypes could be clearly differentiated, all alleles were analysed (H1 and H3). In some cases (M1, M2, M3, and H2), cancer-specific alleles could be detected in the haemocyte samples, but represented less than 50% of the total haemocyte DNA, and the host tissue alleles obscured one or two positions (depending on whether the normal genotype was homozygous or heterozygous at that locus). In these cases, since the tissue allele obscured a potential neoplastic allele, that size was coded as 'missing' for the haemocyte genotype. The total distance between any two samples is calculated as the average of the pairwise differences at all allele sizes observed (104 pairwise comparisons if no data are missing). In pairwise comparisons for each allele, two samples either have a distance of 0 (meaning that the particular allele size is either present in both samples or absent in both samples) or a distance of 1 (meaning that the particular allele size is present in one sample but absent in the other). If data are missing for one or both, then that specific comparison is dropped with no contribution to the total distance between the two samples. The genetic distances calculated are based on the alleles observed at the nine loci, so the absolute value


of the scale itself is therefore dependent on the number of observed alleles that are included in the analysis. Source data for the generation of the phylogenetic tree lists all observed alleles coded as present (1), absent (2), or missing (0) for each sample.

As an alternative analysis method, the data were analysed using Bruvo's method as polyploid data with nine loci, using only the individuals that could be confidently identified. In this alternative analysis, the topology of the resulting tree was not different—all nodes with bootstrap values above 50 were maintained. In this analysis, the genetic distance between the representatives of lineage 1 (H3) and lineage 2 (H1) was 0.676, and the distances to the closest normal sample were 0.595 (H3 to N5) and 0.648 (H1 to N1 and N2).

**qPCR.** Species-specific qPCR of the *EF1 $\alpha$*  locus in *P. aureus* and *V. corrugata* was done using FastStart Universal SYBR Green Master Mix (Roche). Primers were designed to amplify the same region, with the 3' end of both forward and reverse primers on sites that differ between the two species (Extended Data Table 2). Standard curves for each primer set were generated using two control plasmids (one containing the *P. aureus* *EF1 $\alpha$*  fragment and *P. aureus* *SLE* fragment, the other containing the fragments from *V. corrugata*). *EF1 $\alpha$*  fragments were amplified using conserved primers (Extended Data Table 2). Fragments were cloned with a Zero Blunt TOPO Kit (Invitrogen), and plasmids were linearized with NotI before qPCR. Standard curve samples ( $10^4$ – $10^2$  copies per reaction) and experimental samples (2.5 ng per reaction) were done in triplicate. No amplification was detected with the *P. aureus* primers on the *V. corrugata* plasmid or with the *V. corrugata* primers on *P. aureus* plasmid (using up to  $10^7$  copies per reaction). The fraction of *V. corrugata* in each sample was calculated as the copy number using *V. corrugata*-specific primers divided by copy number of *P. aureus*- plus *V. corrugata*-specific amplification.


Quantification of *Steamer*-like elements in the three species was performed by the same method, using primers in the RT-IN region of the retrotransposon (on the basis of sequence obtained through amplification using degenerate primers) and control primers amplifying a region in *EF1 $\alpha$* . All primers used to amplify *SLEs* and *EF1 $\alpha$*  from genomic DNA for cloning into standards and primers for qPCR are listed in Extended Data Table 2. For each species, a single control plasmid containing both the *EF1 $\alpha$*  and *SLE* fragments was used.

30. Muttray, A. F., Schulte, P. M. & Baldwin, S. A. Invertebrate p53-like mRNA isoforms are differentially expressed in mussel haemic neoplasia. *Mar. Environ. Res.* **66**, 412–421 (2008).
31. Howard, D., Lewis, E., Keller, B. & Smith, C. NOAA Technical Memorandum NOS NCCOS Vol. 5 (NOAA/National Centers for Coastal Ocean Science, 2004).
32. Díaz, S., Cao, A., Villalba, A. & Carballal, M. J. Expression of mutant protein p53 and Hsp70 and Hsp90 chaperones in cockles *Cerastoderma edule* affected by neoplasia. *Dis. Aquat. Organ.* **90**, 215–222 (2010).
33. Folmer, O., Black, M., Hoeh, W., Lutz, R. & Vrijenhoek, R. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Mol. Mar. Biol. Biotechnol.* **3**, 294–299 (1994).
34. Oliverio, M. & Mariottini, P. Contrasting morphological and molecular variation in *Coralliophila meyendorffii* (Muricidae, Coralliophilinae). *J. Molluscan Stud.* **67**, 243–245 (2001).
35. Salvi, D. & Mariottini, P. Molecular phylogenetics in 2D: ITS2 rRNA evolution and sequence-structure barcode from Veneridae to Bivalvia. *Mol. Phylogenet. Evol.* **65**, 792–798 (2012).
36. Guindon, S. *et al.* New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst. Biol.* **59**, 307–321 (2010).



**Extended Data Figure 1 | Analysis of *mtCOI* amplified from tissue and haemocyte DNA of normal and diseased mussels (*M. trossulus*).** A partial region of the *mtCOI* gene was amplified from genomic DNA of solid tissue and haemocytes from mussels (*M. trossulus*) and directly sequenced (Fig. 1b). Trace images show a region flanking a representative SNP marked with an open triangle (tissue) or closed triangle

(haemocytes). **a, b**, In normal mussels, tissue and haemocyte alleles match (with G496 at all positions). **c–e**, In mussels with disseminated neoplasia, the tissue and haemocyte alleles are different. Neoplastic haemocytes have A at position 496 and G in tissue, with some A observable in tissue, probably because of infiltration of neoplastic haemocytes.



**Extended Data Figure 2 | Quantification of Steamer-like element genomic copy number in mussels (*M. trossulus*), cockles (*C. edule*), and golden carpet shell clams (*P. aureus*). a–d, Fragments from the SLE reverse transcriptase region and *EF1 $\alpha$*  genes were cloned from each species. Haploid copy numbers of Steamer-like elements (SLE) were quantified by determining the ratio of SLE/*EF1 $\alpha$*  in genomic DNA from haemocytes. Single species-specific SLEs were analysed in (a) mussels**

(*M. trossulus*) and (b) cockles (*C. edule*). c, d, In golden carpet shell clams, one SLE (*SLE-Pa*) was cloned from a normal *P. aureus* (clam N2) and a different one (*SLE-Vc*) was cloned from neoplastic cells (clam H2). Both SLEs could be found in both species, and qPCR analysis confirmed that *SLE-Pa* is more highly amplified in *P. aureus* and has fewer copies in *V. corrugata* and in the neoplastic cells derived from *V. corrugata*.

Extended Data Table 1 | Morphometric analysis of type A and type B cockle (*C. edule*) neoplasia

Cockle ID	Type	Cells counted (N)	Cell diameter ± SD (µm) <sup>*</sup>	Nuclear diameter ± SD (µm) <sup>†</sup>	Nuclear/cell ratio ± SD <sup>‡</sup>
M1	A	15	9.0 ± 1.2	6.7 ± 1.0	0.75 ± 0.09
H2	A	15	8.7 ± 0.9	6.6 ± 0.8	0.76 ± 0.06
H3	A	10	8.5 ± 1.1	6.5 ± 1.0	0.77 ± 0.07
total	A	40	8.8 ± 1.0	6.6 ± 0.9	0.76 ± 0.07
M2	B	10	7.7 ± 0.7	5.5 ± 0.5	0.72 ± 0.07
M3	B	15	6.7 ± 1.2	4.5 ± 0.6	0.70 ± 0.15
H1	B	15	7.3 ± 1.0	5.4 ± 0.8	0.75 ± 0.09
total	B	40	7.2 ± 1.1	5.1 ± 0.8	0.72 ± 0.11

<sup>\*</sup>Cell diameter is statistically different between types A and B ( $P < 0.0001$ ). All tests are two-tailed *t*-tests.

<sup>†</sup>Nuclear diameter is statistically different between types A and B ( $P < 0.0001$ ); for all pairwise comparisons between type A and type B individuals,  $P < 0.05$ .

<sup>‡</sup>Ratio of nuclear diameter to cell diameter is not significantly different between types.

Extended Data Table 2 | Primers used in PCR and qPCR

Target	Forward Primer		Reverse Primer		Temp
<b>mtCOI</b>	LCO1490	GGTCAACAAATCATAAAGATATTGG	HCO2198	TAAACTTCAGGGTGACCAAAAAATCA	50°C
<b>ITS</b>	its-3d	gcgtcgatgaagagcgca	its-4r	agtttttttctcccctta	55°C
<b>EF1<math>\alpha</math></b>					
Mussel ( <i>M. trossulus</i> )	consEF1-F1	ACCATTGATATTGCTYTNITGGAA	MtEF1-R2	ACAATCAAAATGGCACAATC	50°C
Cockle ( <i>C. edule</i> )	CeEF1-F2	TGGTATCACCATCGATATTGC	CeEF1-R3	CCGTTTCGGATCTCTACAGG	55°C
<i>P. aureus</i> & <i>V. corrugata</i>	consVEF1F3b	AGGAACCTCTCAAGCTGAYTG	consVEF1R3b	AGCAATTACCTCGCTGTANGG	55°C
<b>Cockle (<i>C. edule</i>) Microsatellite Loci</b>					
	CeATC1-5F-6FAM	cgttctaccggcatatgtcac	CeATC1-5R	caccttccaccactagaagaaaa	60°C
	CeATC1-22F-VIC	caaacctgaccgggtttatt	CeATC1-22R	tgactccaacttttccagttcca	60°C
	CeATC1-36F-NED	gacatgacaaacaggcctca	CeATC1-36R	tggtcctggtcttatttccac	60°C
	CeATC1-52F-PET	aatctgattttgccacctct	CeATC1-52R	agctcataggagttgtatacgtaa	55°C
	CeATC1-54F-6FAM	tacaaggccgagaaactgct	CeATC1-54R	caatgactgcctcaatgagga	55°C
	CeATC2-4F-VIC	tggaaatgcattcattgagc	CeATC2-4R	cggattgcttctttgatct	55°C
	CeATC2-11F-NED	tgggtgtcaattagatgcttg	CeATC2-11R	taggctcgcaagaaagatggt	60°C
	CeATC2-34F-6FAM	gccatagaggccacctatt	CeATC2-34R	gggctgacaagatttgacatt	60°C
	CeATC2-46F-NED	accaaggcagatatcgtacc	CeATC2-46R	tccagttttaaaccgactctga	55°C
<b>Cloning qPCR Standards</b>					
<b>EF1<math>\alpha</math></b>					
Mussel ( <i>M. trossulus</i> )	consEF1F1	ACCATTGATATTGCTYTNITGGAA	consEF1R2	ACGTTGAAACCAACRTRTC	50°C
Cockle ( <i>C. edule</i> )	CeEF1-F1	TGGAACAACTGAAGCCCGA	CeEF1-R3	CCGTTTCGGATCTCTACAGG	50°C
<i>P. aureus</i> & <i>V. corrugata</i>	consVEF1F3b	AGGAACCTCTCAAGCTGAYTG	consVEF1R3b	AGCAATTACCTCGCTGTANGG	55°C
<b>SLE</b>					
Mussel ( <i>M. trossulus</i> )	DHKPL-F1	TTGAAAGCGACCACAARCCNYT	PXRPW-R1	TTCCGACTTTGGCCCANGGNC	55°C
Cockle ( <i>C. edule</i> )	LWV-F1	GGTATCCAGCCTAGTNGTNGT	PXRPW-R1	TTCCGACTTTGGCCCANGGNC	55°C
<i>P. aureus</i> & <i>V. corrugata</i>	DHKPL-F1	TTGAAAGCGACCACAARCCNYT	PXRPW-VaR1	CTGCAATTTCTGCCANGGNC	55°C
<b>qPCR</b>	<b>Forward Primer</b>		<b>Reverse Primer</b>		<b>Control Plasmids</b>
<b>EF1<math>\alpha</math></b>					
Mussel ( <i>M. trossulus</i> )	MtEF1qF2	TAGGTATTGGAAACAGTGCCAGT	MtEF1qR1	AGACTCGTGGTGCATTTCTAC	pMEF1-SLE4
Cockle ( <i>C. edule</i> )	CeEF1qF2	CCAGTGGGCCGAGTTGAGAC	CeEF1qR2	CTTCAGTGGTCACGTTGGCAG	pCockleSLE-EF1
<i>P. aureus</i>	VaEF1qF1	AAAGAATGGACAGACCAGAGAG	VaEF1qR1	TGCTTCACACCAATGTGAAA	pVaN2-EF1SLE
<i>V. corrugata</i>	VcEF1qF1	CAAGAATGGACAGACAAGAGAA	VcEF1qR1	TGTTTCACACCAAGGTGAAG	pVaH2-EF1SLE
<b>SLE</b>					
<i>SLE-Mt</i>	MtSLEqF2	TGGACTACTTTACAAAAGCCAACG	MtSLEqR1	GCTTCGTGCAATTTCACTAACAT	pMEF1-SLE4
<i>SLE-Ce</i>	cockleRT-F2	GCAGTGTGCAATGTCCAGT	cockleRT-R2	GACCTGTGTCGGAGGCATCA	pCockleSLE-EF1
<i>SLE-Pa</i>	VaNSLEqF1	TAGAGACGAATTTGCTCCGTGG	VaNSLEqR4	CGTGCTCGTGTAAGCATTT	pVaN2-EF1SLE
<i>SLE-Vc</i>	VaHSLEqF1	TAGAGATGAACTCTCTTTGA	VaHSLEqR1	TGCACGGCTTAATGTTTGGGA	pVaH2-EF1SLE